

FORMING THE DUSTY RING IN HR 4796A

SCOTT J. KENYON AND KENNETH WOOD

Smithsonian Astrophysical Observatory, 60 Garden Street, Cambridge, MA 02138

skenyon@cfa.harvard.edu, kwood@cfa.harvard.edu

AND

BARBARA A. WHITNEY AND MICHAEL J. WOLFF

Space Science Institute, Suite 23, 1540 30th Street, Boulder, CO 80303-1012

whitney@colorado.edu, wolff@colorado.edu

Received: 7 July 1999, Accepted: 11 August 1999

ABSTRACT

We describe planetesimal accretion calculations for the dusty ring observed in the nearby A0 star HR 4796A. Models with initial masses of 10–20 times the minimum mass solar nebula produce a ring of width 7–15 AU and height 0.3–0.6 AU at 70 AU in ~ 10 Myr. The ring has a radial optical depth ~ 1 . These results agree with limits derived from infrared images and from the excess infrared luminosity.

Subject headings: planetary systems – solar system: formation – stars: formation – stars: individual (HR 4796A) – circumstellar matter

1. INTRODUCTION

HR 4796A is a nearby A star with a large infrared (IR) excess. Jura (1991) measured the far-IR excess of this wide binary using *IRAS* data. Jura *et al.* (1995, 1998) associated the excess with the A0 primary and derived the ratio of the far-IR to stellar luminosity, $L_{FIR}/L_{\star} \approx 5 \times 10^{-3}$. In 1998, two groups reported extended thermal emission at $\lambda = 20 \mu\text{m}$ from a dusty disk with an inner hole at ~ 40 –70 AU assuming the *Hipparcos* distance of 67 ± 3.5 pc (Jayawardhana *et al.* 1998; Koerner *et al.* 1998). Observations with NICMOS aboard *HST* have revealed a thin annulus of scattered light, with a width of $\lesssim 17$ AU at a distance of ~ 70 AU from the central star (Schneider *et al.* 1999). With an age of ~ 10 Myr (Stauffer *et al.* 1995; Barrado y Navascues *et al.* 1997), the A0 star is older than most pre-main-sequence stars and younger than stars like β Pic and α Lyr with ‘debris disks’.

The dusty ring in HR 4796A challenges theories of planet formation. In most planetesimal accretion calculations, planet-sized objects do not form on short timescales at large distances from the central star. Kenyon & Luu (1999; KL99 hereafter) estimate formation times of 10–40 Myr for Pluto at 35 AU from the Sun. Achieving shorter timescales at 70 AU in HR 4796A requires large initial masses, which might conflict with masses derived from IR observations. In the inner Solar System, planet formation cannot be confined to a narrow ring, because high velocity objects in adjacent annuli interact and ‘mix’ planetary growth over a large area (Weidenschilling *et al.* 1997). This problem may be reduced at larger distances from the central star, where planetary growth is ‘calmer’.

Our goal in this paper is to develop planetesimal accretion models that can lead to the dusty ring observed in HR 4796A. We begin in §2 with Monte Carlo calculations to constrain the geometry and optical depth of dust in the ring. In §3, we derive plausible initial conditions which produce the observed dust distribution on 10 Myr timescales. These models also satisfy constraints

on the dust mass from *IRAS* observations and lead to a self-consistent picture for ring formation. We conclude in §4 with a brief summary and discussion of the implications of this study for planet formation in other star systems.

2. MODEL IMAGES

Current data constrain the geometry and optical depth of the ring. Near-IR images measure the amount of scattered light from the ratio of the 1.1–1.6 μm radiation to the stellar luminosity, $L_{NIR}/L_{\star} \approx 2 \times 10^{-3}$ (Schneider *et al.* 1999). The far-IR luminosity limits the amount of stellar radiation absorbed and reradiated. To construct a physical model, we assume an annulus of width Δa and height z at a distance $a = 70$ AU from the central star. The luminosity ratios depend on the solid angle $\Omega/4\pi = 2\pi az/4\pi a^2 = z/2a$, the radial optical depth τ , and the albedo ω : $L_{NIR}/L_{\star} = \tau\omega(z/2a)$ and $L_{FIR}/L_{\star} = \tau(1 - \omega)(z/2a)$. These equations assume gray opacity and scattering in the geometric optics limit. If the annulus contains planetesimals and dust in dynamical equilibrium, $z/\Delta a \lesssim 1$ (Hornung *et al.* 1985). Anticipating the results of our coagulation calculations, where $z/a \sim 10^{-2}$, we then have $\omega \approx 0.3$ – close to observed values in β Pic (Backman & Paresce 1993) – and $\tau \sim 1$.

We construct scattered light images using a 3D Monte Carlo code (Wood & Reynolds 1999) with forced first scattering (Witt 1977) and a ‘peeling-off’ procedure (Yusef-Zadeh, Morris, & White 1984). We adopt a dust number density, $n = n_0 e^{-z^2/2H^2} e^{-(a-70)^2/2A^2}$, where the scale height H and scale length A are in AU. We assume $\omega = 0.3$, isotropic scattering (see Figure 12 of Augereau *et al.* 1999), and adjust τ until the models yield $L_{NIR}/L_{\star} = 1.5 \times 10^{-3}$ for an input H and A . Model images with $\omega\tau = \text{constant}$ are identical in the optically thin limit.

Figure 1 compares several models with the NICMOS 1.1 μm image (from FITS data kindly sent by G. Schneider). We convolved Monte Carlo images with a gaussian point-spread function with FWHM = $0''.12$ to approximate

the $0''.12$ resolution of NICMOS (Schneider *et al.* 1999). Model images with $H > 5$ AU (FWHM = 14 AU) or $A > 10$ AU (FWHM = 27 AU) are more extended than the data (Augereau *et al.* 1999). Our preferred model with $H = 0.5$ AU, $A = 5$ AU, and $\omega\tau = 0.25$ reproduces the size and shape of the NICMOS image as well as the limb brightening observed towards the ring edges. These results match NICMOS flux ratios best for our adopted geometry; larger H implies smaller $\omega\tau$. The 3σ limit, $\omega\tau = 0.12$ – 0.35 , agrees with previous estimates (cf. Koerner *et al.* 1998; Schneider *et al.* 1999; Augereau *et al.* 1999). We disagree, however, with the $\tau \sim 10^{-3}$ of Schneider *et al.*; their result is valid only for scattering in a spherical shell.

3. COAGULATION MODEL

To calculate dust evolution in HR 4976A, we use a coagulation code based on the particle-in-a-box method (KL99). This formalism treats planetesimals as a statistical ensemble of bodies with a distribution of horizontal and vertical velocities about Keplerian orbits (Safronov 1969). We begin with a size distribution of N_i bodies having total mass M_i in each of i mass batches. Collisions among these bodies produce (i) growth through mergers along with cratering debris for low impact velocities or (ii) catastrophic disruption into numerous small fragments for high impact velocities. Inelastic collisions, long range gravitational interactions (dynamical friction and viscous stirring), and gas drag change the velocities of the mass batches with time. The code has been tested against analytic solutions of the coagulation equation and published calculations of planetesimal growth. Although inappropriate for the last stages of planet formation, our approach well-approximates the early stages (Kokubo & Ida 1996).

We model planetesimal growth in an annulus of width $\Delta a = 12$ AU centered at $a = 70$ AU. The central star has a mass of $2.5 M_\odot$. The input size distribution has equal mass in each of 38 mass batches with initial radii $r_i = 1$ – 80 m. For a Minimum Mass Solar Nebula with mass M_{MMSN} , the total mass in the annulus is $M_0 \approx 15 M_E$; the initial number of bodies with $r_i = 1$ m is $N_0 \approx 3 \times 10^{20}$. All batches start with the same initial velocity. The mass density $\rho_0 = 1.5 \text{ g cm}^{-3}$, intrinsic strength $S_0 = 2 \times 10^6 \text{ erg g}^{-1}$, and other bulk properties of the grains are adopted from earlier work (see KL99).

Planetesimal growth at 70 AU follows the evolution described previously (KL99). The 80 m bodies first grow slowly into 1 km objects. During this slow growth phase, frequent collisions damp the velocity dispersion of all bodies. “Runaway growth” begins when the gravitational range of large objects exceeds their geometric cross-section. These bodies grow from 1 km up to ~ 100 km in several Myr. During runaway growth, collisional debris, dynamical friction, and viscous stirring increase the velocity dispersion of small bodies from $\sim 1 \text{ m s}^{-1}$ up to $\sim 40 \text{ m s}^{-1}$. This evolution reduces the gravitational range of the 100 km objects and ends runaway growth. The largest objects then grow slowly to 1000+ km sizes.

Figure 2(a) shows the growth of the largest object in several models. For $M_0 = 10 M_{MMSN}$ and $e_0 = 10^{-3}$, Pluto-sized objects form in $t_P = 2.1$ Myr at $a = 35$ AU, 13 Myr at 70 AU, and 93 Myr at 140 AU. Models with smaller M_0 take longer to make “Pluto”. Plutos form more quickly for $e_0 < 10^{-3}$, because gravitational focusing factors are

larger.

Figure 2(b) shows the evolution of the scale height H for small objects. Initially, $H = 2\pi a \sin i \leq 0.003a$ for $e_0 \leq 10^{-3}$. Collisional damping cools the bodies during the slow-growth phase; H remains small. H increases dramatically during runaway growth, when dynamical processes heat up the smallest bodies. Once runaway growth ends, H slowly increases to 0.3–0.6 AU independent of M_0 , e_0 , and other input parameters.

When H begins to increase, high velocity collisions produce numerous “dust grains” with sizes $\lesssim 1$ m. We do not follow explicitly the evolution of these bodies. Instead, we assume that collisional debris is (i) swept up by 1 m or larger objects, (ii) ejected by radiation pressure, or (iii) dragged inwards by the Poynting-Robertson effect. Grains with sizes exceeding 4–5 μm are stable against radiation pressure (Jura *et al.* 1998; Augereau *et al.* 1999). Poynting-Robertson drag reduces the mass in small grains on a timescale $t_{PR} \approx 1.0$ Myr ($r_i/4 \mu\text{m}$). With the short collision times, $\leq 10^5$ yr, in our model annulus, 1 Myr seems a reasonable estimate of the timescale for collisions to produce 4 μm grains which are removed by radiative processes. For this paper, we calculate the accretion explicitly and adopt a 1 Myr timescale for dust removal.

Figure 2(c) shows the dust mass as a function of time. The results are not sensitive to the adopted mass distribution for grains with $r_i \gtrsim 4 \mu\text{m}$ or to factor of 2–3 variations in the removal timescale. The dust mass is initially large due to the starting conditions. The dust mass decreases with time, because (i) collisional damping of the smaller bodies leads to less collisional debris and (ii) radiative processes and accretion by large bodies remove dust. Once runaway growth begins, collisions between small bodies produce more dust. The dust mass then reaches a rough equilibrium between collision debris and dust removed by radiation forces and by the larger bodies.

These results indicate that large dust masses correlate with runaway growth and the formation of 1 or more Plutos in the outer parts of the disk. To predict the amount of radiation absorbed and scattered by dust and larger bodies, we compute τ from the model size distribution. We assume the geometric optics limit because $r_i \gg \lambda$. For the large bodies $\tau = \sum_{i=1}^N n_i \sigma_i \Delta a$, where n_i is the number density in mass batch i , σ_i is the extinction cross-section, and N is the number of mass batches. We adopt $\sigma_i = 2\pi r_i^2$ and a volume $V_i = 2\pi a \Delta a H_i$ to compute $n_i = N_i/V_i$ and hence τ for material with $r_i \gtrsim 1$ m.

Estimating τ for small particles requires an adopted cumulative size distribution, $N_C \propto r_i^{-q}$. We consider three choices: (i) $q = 2.5$, the collisional limit for coagulation; (ii) $q = 3$, equal mass per mass interval; and (iii) $q = 3.5$, the approximate distribution for grains in the interstellar medium. Our calculations produce $q \approx 2.7$ for 1–100 m bodies. We expect a slightly steeper mass distribution for smaller bodies, because collisions between smaller bodies produce fewer mergers and more debris.

Figure 2(d) shows how τ evolves for a single model. The large bodies initially have modest radial optical depth, $\tau_L \approx 0.2$. This optical depth decreases with time, except for a brief period when runaway growth produces 10–100 km objects with small scale height above the disk midplane. The large bodies are transparent once a Pluto

forms. The small grains are also initially opaque. This dust is transparent at late times if most of the mass is in the largest grains, $q \lesssim 2.8$. The dust is opaque for $q \gtrsim 3$.

Table 1 summarizes results for various initial conditions. Models with $M_0 \approx 10\text{--}20 M_{MMSN}$ and $e_0 \approx 10^{-4}\text{--}10^{-3}$ achieve $\tau \sim 1$ in 10 Myr. Less massive disks produce less dust on longer timescales. The results are not sensitive to other input parameters, including the size distribution and the bulk properties of the bodies.

Table 1 also shows why dust in HR 4796A lies in a ring. In disks with surface density $\Sigma \propto a^{-3/2}$, the Pluto formation timescale¹ is $t_P \approx 13 \text{ Myr } (M_0/10 M_{MMSN})^{-1} (a/70 \text{ AU})^{2.7}$. Once an annulus at a begins to form dust, material at $a + \Delta a$ must wait a time, $\Delta t/t_P \approx 2.7\Delta a/a$, to reach the same state. This result sets a hard outer limit to the ring, $\Delta a/a \approx 0.4\Delta t/t_P \approx 0.1\text{--}0.2$, if Δt is the time for H to double in size during runaway growth, 2–3 Myr. We expect a hard inner edge, because particle velocities reach the shattering limit of $\sim 100 \text{ m s}^{-1}$ (KL99) or planets sweep up the dust (e.g., Pollack *et al.* 1996) or both.

4. DISCUSSION AND SUMMARY

Our results indicate that the dusty ring in HR 4796A is a natural outcome of planetesimal evolution. Planet formation at 70 AU in 10 Myr is possible with an initial disk mass of $10\text{--}20 M_{MMSN}$. Dust production associated with planet formation is then confined to a ring with $\Delta a \approx 7\text{--}15 \text{ AU}$. The optical depth in this ring satisfies current constraints on scattered light at $1\text{--}2 \mu\text{m}$ and on thermal emission at $10\text{--}100 \mu\text{m}$ if the size distribution of the dust is $N_C \propto r_i^{-q}$ with $q \gtrsim 3$ for $r_i \lesssim 1 \text{ m}$. Models with disk masses smaller than $10 M_{MMSN}$ fail to produce planets and an observable dusty ring in 10 Myr.

An uncertainty in our model is the timescale to produce $1\text{--}80 \text{ m}$ bodies from small dust grains in a turbulent, gaseous disk. Cuzzi *et al.* (1993) show that grains grow very rapidly once they decouple from eddies in the disk. The decoupling timescale depends on the unknown disk viscosity at 70 AU.

Our model makes several observational predictions. We expect $L_{NIR}/L_\star = \text{constant}$ for $\lambda \leq 5 \mu\text{m}$; current data are consistent with this prediction at the 1.5σ level. Bet-

ter measurements of the ring flux at $\lambda \geq 1.6 \mu\text{m}$ would test our optical depth assumptions and yield interesting constraints on grain properties. Deep images at $\lambda \geq 10\text{--}20 \mu\text{m}$ with high spatial resolution should detect material outside the ring. We predict $\tau \approx 0.1$ in large bodies for $a \gtrsim 80 \text{ AU}$; the surface brightness and temperature of this material should decrease markedly with radius. This material should have negligible mass in small objects, because coagulation concentrates mass in the largest objects when H is small. We also expect a flux of dust grains into the central star, although we cannot yet compare quantitative predictions with observations. Future calculations of radiative processes within the ring will address this issue.

Applying this HR 4796A model to other stars with circumstellar disks is challenging due to small number statistics and unfavorable circumstances. Nearby companion stars probably influence the dynamics of dusty rings in HD 98800 and HD 141569 (Pirzkal *et al.* 1997; Low *et al.* 1999; Lagrange *et al.* 2000). In HR 4796A, the M-type companion lies well outside the ring radius and cannot modify ring dynamics significantly. Older systems like β Pic and α Lyr require time-dependent treatment of dust to allow the ring to spread with time (e.g., Artymowicz 1997). We plan to incorporate this time-dependent behavior in future calculations to see whether the ring in HR 4796A can evolve into a debris disk (as in e.g. β Pic and α Lyr) on a timescale of $100\text{--}200 \text{ Myr}$.

The main alternative to *in situ* ring formation at 70 AU is migration of a planet formed at a smaller radius. Weidenschilling & Marzari (1996) show that gravitational interactions can scatter large objects into the outer disk in less than 1 Myr. Migration reduces the required ring mass by a factor of $10\text{--}100$. However, the scattered body has a large eccentricity, $e \sim 0.5$. Dynamical friction might circularize the orbit in 10 Myr, but would induce large eccentricities in smaller bodies. The width of the dusty ring would probably exceed observational constraints. Future calculations can address these issues.

We thank B. Bromley for helping us run our code on the HP Exemplar “Neptune” at JPL and for a generous allotment of computer time through funding from the NASA Offices of Mission to Planet Earth, Aeronautics, and Space Science.

REFERENCES

- Artymowicz, P., Burrows, C., & Paresce, F. 1989, *ApJ*, 337, 494
 Artymowicz, P. 1997, *ARE&PS*, 25, 175
 Augereau, J. C., Lagrange, A.-M., Mouillet, D., Papaloizou, J. C. B., & Grorod, P. A. 1999, *A&A*, in press (astro-ph/9906429)
 Backman, D. E., & Paresce, F. 1993, in *Protostars and Planets III*, eds. E. H. Levy & J. I. Lunine, Tucson, Univ of Arizona, p. 1253
 Barrado y Navascues, D., Stauffer, J. R., Hartmann, L., & Balachandran, S. C. 1997, *ApJ*, 475, 313
 Cuzzi, J. N., Dobrovolskis, A. R., & Champney, J. M. 1993, *Icarus*, 106, 102
 Hornung, P., Pellat, R., & Barge, P. 1985 *Icarus*, 64, 295
 Jayawardhana, R. *et al.* 1998, *ApJ*, 503, L79
 Jura, M. 1991, *ApJ*, 383, L79
 Jura, M., Ghez, A. M., White, R. J., McCarthy, D. W., Smith, R. C., & Martin, P. G. 1995, *ApJ*, 445, 451
 Jura, M., Malkan, M., White, R., Telesco, C., Pina, R., & Fisher, R. S. 1998, *ApJ*, 505, 897
 Kenyon, S. J., & Luu, J. X. 1999, *AJ*, 118, No. 2 (astro-ph/9904115)
 Koerner, D. W., Ressler, M. E., Werner, M. W., & Backman, D. E. 1998, *ApJ*, 503, L83
 Kokubo, E., & Ida, S. 1996, *Icarus*, 123, 180
 Lagrange, A.-M., Backman, D., & Artymowicz, P. 2000, in *Protostars & Planets IV*, eds. V. Mannings, A. P. Boss, & S. S. Russell, Tucson, Univ. of Arizona, in press
 Low, F. J., Hines, D. C., & Schneider, G. 1999, astro-ph/9905366
 Pirzkal, N., Spillar, E. J., & Dyck, H. M. 1997, *ApJ*, 481, 392
 Pollack, J. B., Hubickyj, O., Bodenheimer, P., Lissauer, J. J., Podolak, M., & Greenzweig, Y. 1996, *Icarus*, 124, 62
 Safronov, V. S. 1969, *Evolution of the Protoplanetary Cloud and Formation of the Earth and Planets*, Nauka, Moscow [Translation 1972, NASA TT F-677]
 Schneider, G., *et al.* 1999, *ApJ*, 513, L127
 Stauffer, J. R., Hartmann, L. W., & Barrado y Navascues, D. 1995, *ApJ*, 454, 910
 Weidenschilling, S. J., Spaute, D., Davis, D. R., Marzari, F., Ohtsuki, K. 1997, *Icarus*, 128, 429
 Weidenschilling, S. J., & Marzari, F. 1996, *Nature*, 384, 619
 Witt, A.N. 1977, *ApJS*, 35, 1
 Wood, K., & Reynolds, R.J. 1999, *ApJ*, in press, astro-ph/9905289
 Yusef-Zadeh, F., Morris, M., & White, R.L. 1984, *ApJ*, 278, 186

¹Pluto is a handy reference: $1000+$ km objects form roughly in the middle of the rapid increase in H which produces large dust masses.




FIG. 1.— Comparison of model scattered light images with HST data of HR 4796A. (a) upper left panel: NICMOS coronagraphic image at $1.1 \mu\text{m}$. (b) upper right panel: Model scattered light image with $z = 0.5 \text{ AU}$, $R = 5 \text{ AU}$, and $\omega\tau_{NIR} = 0.25$. (c) lower left panel: As in (b) for $z = 5 \text{ AU}$, $R = 10 \text{ AU}$, and $\omega\tau_{NIR} = 0.02$. (d) lower right panel: As in (c) for $z = 1 \text{ AU}$, $R = 20 \text{ AU}$, and $\omega\tau_{NIR} = 0.1$.




FIG. 2.— Results for HR 4796A coagulation models. (a) lower left panel: maximum radius for models with $e_0 = 10^{-3}$ at $a = 35, 70,$ and 140 AU; initial masses are listed in the legend. (b) upper left panel: scale height of small bodies for 70 AU models with e_0 and M_0 as listed. (c) upper right panel: dust mass for 70 AU models with e_0 and M_0 as listed. (d) lower right panel: optical depth for a 70 AU model with $e_0 = 10^{-3}$ and $M_0 = 10M_{MMSN}$.

TABLE 1
RESULTS OF PLANETESIMAL ACCRETION CALCULATIONS

a	M_0	e_0	t_P	$\log \tau_S$	$\log \tau_L$
35	100	10^{-3}	2.1	0.09	-3.43
70	15	10^{-4}	81.1	-1.90	-3.92
70	45	10^{-4}	20.4	-0.96	-3.27
70	150	10^{-4}	5.7	-0.04	-2.69
70	15	10^{-3}	156.4	-2.21	-4.24
70	45	10^{-3}	50.0	-1.42	-3.65
70	75	10^{-3}	29.9	-0.94	-3.48
70	150	10^{-3}	13.0	-0.34	-3.25
70	300	10^{-3}	6.6	-0.05	-2.62
140	200	10^{-3}	92.6	-1.20	-2.95

NOTE.— a is the distance of the annulus from the central star in AU; M_0 is the initial mass in the annulus in M_E ; e_0 is the initial eccentricity of each mass batch; t_P is the timescale in Myr to produce Pluto-sized objects; τ_S is the optical depth in dust when the first Pluto forms, assuming equal mass in dust per decade in radius; τ_L is the optical depth of the large bodies when the first Pluto forms.